

- Morris, R.F. (editor). 1963. The dynamics of epidemic spruce budworm populations. *Memoirs Ent. Soc. Can.* No. 31, 332 pp.
- Schmitt, D.M., Grimble, D.G. and Searcy, J.L. 1984. Managing the Spruce Budworm in Eastern North America. U.S.D.A. Forest Service. Agriculture Handbook No. 620.
- Stokes, M.A. and T.L. Smiley. 1968. An introduction to tree-ring dating. U. of Chicago Press, Chicago 73 pp.
- Thomson, A.J., R.F. Shepherd, J.W.E. Harris and R.H. Silversides. 1984. Relating weather to outbreaks of western spruce budworm *Choristoneura occidentalis* (Lepidoptera: Tortricidae) in British Columbia. *Can. Ent.* 116: 375-381.
- Wellington, W.R., J.J. Fettes, K.B. Turner and R.M. Belyea. 1950. Physical and biological indicators of the development of outbreaks of the spruce budworm. *Can. J. Research* 28: 308-331.

***CARNUS HEMAPTERUS* (DIPTERA: CARNIDAE) AN AVIAN NEST PARASITE NEW TO BRITISH COLUMBIA**

RICHARD J. CANNINGS

Cowan Vertebrate Museum
Department of Zoology
University of British Columbia
Vancouver, B.C. V6T 2A9

Carnus hemapterus Nitzsch is an ectoparasite of bird nestlings found throughout Europe and scattered locations in North America. (Bequaert 1942, Capelle and Whitworth 1973). Although Sabrosky (1965) lists it only from New Brunswick in Canada, Bequaert (1951) did state that *C. hemapterus* was also "found in British Columbia...the details...to be published later by the discoverers." As far as I know, those details were never published.

While checking a nest of the Northern Saw-whet Owl (*Aegolius acadicus*) near Osoyoos, B.C. on April 17, 1985, I noticed several small flies crawling over the newly-hatched nestlings. I collected a few specimens on

April 17, 19, and 21, and on April 25 I took 50 flies off two nestlings. They were identified as *C. hemapterus* by S.G. Cannings and J.F. McAlpine; voucher specimens are now at the University of British Columbia, Canadian National Collection, and the University of Guelph.

C. hemapterus has been collected from the nests of a wide variety of birds, but primarily from those of raptors and hole-nesting species. I found it to be common in the Osoyoos area, being present in all of 13 nests of the European Starling *Sturnus vulgaris* and two other Northern Saw-whet Owl nests that I checked. Further details of the infestations are being published elsewhere (Cannings, in press).

REFERENCES

- Bequaert, J. 1942. *Carnus hemapterus* Nitzsch, an ectoparasitic fly of birds, new to America (Diptera). *Bull. Brook Ent. Soc.* 37: 140-149.
- Bequaert, J. 1951. *Carnus hemapterus* Nitzsch on a screech owl in Arizona (Diptera). *Psyche* 58: 157.
- Cannings, R.J. in press. Infestations of *Carnus hemapterus* Nitzsch (Diptera: Carnidae) in Northern Saw-whet Owl nests. *Murrelet*.
- Capelle, K.J. and T.L. Whitworth. 1973. The distribution and avian hosts of *Carnus hemapterus* (Diptera: Milichiidae) in North America. *J. Med. Ent.* 10: 525-526.
- Sabrosky, C. 1965. Milichiidae. P. 728-733 *In* A. Stone, C.W. Sabrosky, W.W. Wirth, B.H. Foote, and J.R. Coulson (Eds.), A catalog of the Diptera of America north of Mexico. USDA Agric. Res. Serv., Washington, DC, 1696 pp.