

AN ANNOTATED LIST OF THE COLEOPTERA TAKEN AT OR NEAR TERRACE, BRITISH COLUMBIA. PART 2¹

M. E. CLARK
Masset, B.C.

ORTHOPTERIDAE

Sacium lugubre LeC.—Vidit H. C. Fall.

STAPHYLINIDAE

Micropeplus costatus LeC.—Vidit H. C. Fall.
Micropeplus sp.—On fungus. Unknown to Fall, and not in his collection.

Emalus nigrella LeC.—Taken under pine bark, from tunnels of bark beetles.

Proteinus basalis LeC.—Near one in the LeConte collection; November 3, 1923.

Proteinus n. sp.—Taken June and July, 1931. Probably now in the Fall collection.

Megarthus sinuaticollis Boisd.—Taken June and July, 1931; vidit H. C. Fall.

Megarthus sp.—Unknown species, taken in rotten leaves, March 19, 1937.

Anthobium pothos Mann.—June 12, 1920; vidit H. C. Fall and E. C. Van Dyke.

Anthobium sp.—An unknown species.

Acutia tumidula Makl.—Vidit H. C. Fall.

Phyllodrepa sp. near *megarthroides* Fauv.; det. H. C. Fall.

Phyllodrepa n. sp.

Omalium ricularae Payk.—June and July, by sifting and in fungus; vidit H. C. Fall.

Omalium humerosum Fauv.?—Vidit H. C. Fall, who said "vicinity of."

Omalium n. sp., near *longulum* Makl.; det. H. C. Fall.

Omalium spp.—Four unknown species.

Phloeonomus lapponicus Zett.

Porrhodites fenestralis Zett.—Vidit H. C. Fall.

Lathrimacum pictum Fauv.

Lathrimacum humerale Csy.—Taken November 3, 1923.

Lathrimacum reflexicolle Csy.—From mouse nest, March 19, 1935; vidit H. C. Fall.

Lathrimacum sp.—An unknown species from Thornhill Mountain; vidit H. C. Fall.

Arpadium sp.—Three specimens, November 3, 1923, and in 1927; unknown to H. C. Fall.

Acidota sp.—An unknown species taken in 1931.

Amphichroum maculatum Horn—On spathes of skunk cabbage.

Pelecomalium testaceum Mann.—June, 1928, on skunk cabbage; vidit E. C. Van Dyke.

Tilca ruftarsis Csy.—From Thornhill Mountain; vidit R. E. Blackwelder.

Ancyrophorus bimpressus Makl.—Taken on November 24, 1923.

Trogophloeus sp.—An unknown species.

Aploderus linearis LeC.—Taken March 15, 1937; det. H. C. Fall.

Oxytelus fuscipennis Mann.

Oxytelus nitidulus Grav., or possibly *suspectus* Csy.—Collected March 18, 1937; det. H. C. Fall.

Platystethus americanus Er.

Bledius mysticus Fall.

Bledius sp.—An unknown species.

Stenus vexatus Csy.—From muddy puddle of drainage on flat.

Stenus insularis Csy.—From moss on flats.

Stenus corvus Csy.—Taken at Lakelse Lake.

Stenus convictor Csy.—Taken at Lakelse Lake, and in moss, November 5, 1923.

Stenus sp., near *convictor*.—Taken February, 1940.

Stenus nanulus Csy.—Taken at Lakelse Lake, March, 1927, and in rotten leaves.

Stenus sp.—Undescribed species; Lakelse Lake.

Stenus maritimus Mots.—In moss, March, 1935.

Stenus pterobrachys G. & H.—Scarce, June, 1931, and February, 1940.

Stenus alpicola Fauv.—In moss, March 21, 1937.

Stenus egenus Er.—Taken in 1927.

Stenus n. sp.—Collected February, 1940.

Stenus n. sp., nearest *curtus* Csy.—Taken in 1933.

Stenus punctiger Csy.—Taken while sifting, August, 1927.

Stenus sectator Csy.

Stenus sp., near *sectator*.—Vidit H. C. Fall.

Stenus mammops Csy.—Taken in June, 1931.

Stenus reconditus Csy.

Stenus rugifer Csy.

Stenus austini Csy.—In moss, March 18, 1935.

Stenus juno Fab.—Two females and one male; June, 1938.

Stenus pollens Csy.—Taken in 1939.

Stenus sp.—A new species near *monticola* Csy.

Euaestethus sp.—"Species seems new"; det. M. W. Sanderson. Taken by sifting, June and July, 1931.

Paederus pugetensis Csy.—Rare.

Lathrobium rigidum Csy.?—Or near *rigidum*; taken in 1933 and 1938.

Lathrobium sp.—Species near *simile* LeC. Taken in 1938.

Tetartopeus finitimus LeC.—Collected June 1, 1920; one sent to Casey.

Lathrotropis sp.—An unknown species.

?*Medon* sp.—"Or near"; March 4.

Orus punctatus Csy.—Vidit H. C. Fall.

Stilicis oregonus Csy.—Rare; taken in 1931.

Astenus longiusculus Man.

Nudobius cephalus Say.—Rare.

Nudobius spp.—Two unknown species.

Gyrohyppus fusciceps LeC.

Gyrohyppus sp. near *fusciceps*.

Gyrohyppus obsidianus Melsh.—Rare, taken in 1931; vidit H. C. Fall.

Gyrohyppus obscurus Er. — Species vera, teste H. C. Fall.

¹Part I appeared in the previous issue of this journal, pp. 24-27. I am again greatly indebted to Mr. C. A. Frost, who has checked the manuscript, and who in past years submitted many of my beetles to the late Dr. H. C. Fall and others for identification or verification.

- Leptacinodes nigrifulus* LeC.—February 6, 1926.
- Hesperolinus piceus* Csy.—One pair taken in 1933.
- Hesperolinus brunnescens* LeC.
- Parothius californicus* Mann.—Vidit H. C. Fall.
- Baptolinus macrocephalus* Nordm.
- Philonthus*—(All species passed on by H. C. Fall).
- Philonthus furvus* Nordm.
- Philonthus* sp., near *politus* Linn.
- Philonthus* sp., near *atratus* Grav.—Under grizzly bear bones.
- Philonthus varians* Payk.
- Philonthus* sp.—Unknown to C. A. Frost and H. C. Fall.
- Philonthus sordidus* Grav.
- Philonthus crotchi* Horn, or near.
- Philonthus nigrifulus* Grav.—Scarce.
- Philonthus nigrifulus* Grav., or near.—In numbers in old sacks, March, 1937.
- Philonthus longicornis* Steph.—Rare.
- Philonthus aurulentus* Horn.
- Philonthus tetragonocephalus* Notman.—Scarce.
- Philonthus agilis* Grav.—Taken in 1931.
- Philonthus punctatellus* Horn. — Taken in 1937.
- Philonthus microphthalmus* Horn. — Taken in 1937.
- Philonthus quadricollis* Horn. — Taken in 1937.
- Philonthus debilis* Grav.—Taken in 1937.
- Philonthus* sp., near *quadricollis*.—Taken in 1937.
- Staphylinus fossator* Grav.—About stables.
- Staphylinus pleuralis* LeC.—On dung, and under boards.
- Staphylinus caesareus* Cederhj.
- Ontholestes cingulatus* Grav.—Frequent, on cow and horse dung.
- Creophilus villosus* Grav.—Scarce; on carrion.
- Heterothops californicus* LeC. — Taken in 1931; vidit H. C. Fall.
- Heterothops carbonatus* Fall.—Vidit H. C. Fall.
- Quedius* sp., near *marginalis* Makl.—Taken in 1931.
- Quedius laevigatus* Gyll.
- Quedius mesomelinus* Marsh.
- Quedius aenescens* Makl., or near — Under grizzly bear bones.
- Tachinus maculicollis* Makl.—From fungus, November, 1919.
- Tachinus basalis* Er.
- Tachinus crotchi* Horn.
- Tachinus circumcinctus* Makl. — Taken in August, 1927.
- Tachinus nigricornis* Mann.—Taken by sifting in 1927 and 1931.
- Tachinus instabilis* Makl.—Taken in 1931.
- Tachinus tachyporoides* Horn. — Taken in 1931.
- Tachyporus acaudus* Say.—Under bits of wood, loose bark, etc., frequent in spring; det. Blackwelder.
- Tachyporus nitidulus* Fab.—Det. Blackwelder.
- Tachyporus rulomus* Blkwr.—Taken March 9, 1927; det. Blackwelder.
- Tachyporus jocosus* Say.?
- Bolitobius cincticollis* Say.
- Bolitobius poecilus* Mann.
- Bolitobius obsoletus* Say., or near.—H. C. Fall remarked "No concussion."
- Bolitobius* n. sp., near *intrusus* Horn.
- Bolitobius intrusus* Horn.
- Bryoporus rufescens* LeC.
- Mycetoporus flavicollis* LeC.—Taken by sifting; June and July, 1931
- Mycetoporus humidus* Say.—In rotting fungus, June and July, 1931.
- Mycetoporus splendidus* Grav.—According to H. C. Fall.
- Deinopsis* spp.—Several unidentified species.
- Myllaena* sp.—Not identified.
- Datomicra zosteræ* Thoms.
- Acrotona fungi* Grav.—Taken on decayed soft fungus.
- Aleochara bimaculata* Grav.
- Aleochara* sp.—Taken in 1938. Species not known to H. C. Fall.

PSELAPHIDAE

- Sonoma corticina* Csy.—Rare; found under a board in April.
- Sonoma parviceps* Makl.—Rare; taken by sifting in June and July; vidit A. S. Nicolay.
- Cupila clavicornis* Makl.—Very rare; det. A. S. Nicolay.
- Actium retractum* Csy.—Two specimens; vidit A. S. Nicolay.
- Actium pacificum* Csy., "or near".—Det. H. C. Fall.
- Reichenbachia* spp.—Two undescribed species; vidit H. C. Fall.
- Tychus tenellus* LeC.—Rare; shaken from moss, November, 1920.
- Tychus cognatus* LeC.—Rare; taken by sifting, June and July, 1931.
- Pselaphus bellax* Csy.—Rare; taken by sifting, June and July, 1931; vidit H. C. Fall.

PTILIIDAE

- Ptenidium pullum* Makl., "or near; legs lighter" (note by H. C. Fall). From a squirrel's nest, February 4 and March 18, 1935.
- Acratrichis* spp.—A large and a small species in moss in November, both unidentified.
- Ptiliidae—A dozen specimens of an unidentified species, from a mouse nest.

SCAPHIDIIDAE

- Scaphium castanipes* Kby.—Taken in 1932.
- Scaphisoma convexum* Say?—Taken under bark.
- Scaphisoma castaneum* Mots.—Vidit H. C. Fall.
- Scaphidiidae—An unidentified species.

SPHAERITIDAE

- Sphaerites politus* Mann.—Rare; taken beside Alwynne Creek, on bleaching bear bones.

HISTERIDAE

- Hister umbilicatus* Csy.—"Not quite typical," T. L. Casey said of one sent to him.

Hister umbrosus Csy.—Found about the stable. Fall called this a synonym of the above.

Hister immunis Er.—On sand near the river, 1934. Vidit H. C. Fall, but called *fidelis* Csy. by C. G. Siepmann.

Isolomalus mancus Csy.—In numbers under spruce bark, August, 1927.

Plegaderus sp.—Unknown species; vidit H. C. Fall. Very rare under hemlock bark on fallen trees from which bark is stripping.

Saprinus estriatus LeC.—Scarce, sand bank on slough near Skeena River, August, 1927.

Saprinus oregonensis LeC.

Saprinus bigemmeus LeC.—Mr. Frost wrote "Not quite like some from California sea coast."

LYCIDAE

Eros simplicipes Mann.—Frequent in the woods; May 25, 1929; June; det. E. C. Van Dyke.

Eros aurora Hbst.—Scarce; in the woods, June.

Eros nigripes Schaeff.—Scarce; in flight, first week of May; det. by C. Schaeffer and E. C. Van Dyke.

Eros thoracicus Rand.—Scarce; vidit H. C. Fall.

Plateros sp.—An unknown species.

Plateros californicus Van Dyke?—Vidit H. C. Fall.

LAMPYRIDAE

Lucidota corrusca Linn.—Common on flowers and herbage in the spring, and on wild everlasting flowers in the fall.

Lucidota lacustris LeC.—One of my specimens was compared with the type.

CANTHARIDAE

Podabrus piniphilus Esch.—Scarce; flying in May, June and July.

Podabrus comes LeC.

Silis pallida Mann.—June 14, 1920.

Malthodes sp.—May have been *humidus* Fender or *oregonus* Fender.

MELASIDAE

Melasis rufipennis Horn.—Rarely found, but was breeding in numbers in a confined spot in punky hemlock.

Isorhipis ruficornis Say.—"New record for the west." Taken in 1933.

Microrhagus pectinatus LeC.—"New record for the west."

Epiphaniis cornutus Esch.

THROSCIDAE

Pactopus hornii LeC.—Scarce; flying; under moss in November, under cover in June; vidit H. C. Fall.

Throscus carinicoellis Schaeff.—Det. H. C. Fall.

HETERO CERIDAE

Heteroceris brunneus Melsh.—H. C. Fall's note: "runs to, but not the true one."

DASCILLIDAE

Macropogon piceus Horn?—"May be *dubius* Brown." Scarce; under dry moss flakes in May; on rocks of Little Canyon, Terrace.

Araeopidius monachus LeC.

Eucinetus terminalis LeC.—"New record for the west"; det. H. C. Fall, 1937.

Eucinetus sp.—An undescribed species; in Mr. C. A. Frost's collection.

HELODIDAE

Cyphon variabilis Thunb.

Cyphon concinnus LeC.

BYTURIDAE

Byturus bakeri Barber?—Vidit H. S. Barber; a bad pest on the wild blackcaps and raspberries.

DERMESTIDAE

Dermestes lardarius Linn.—A common pest; also taken on Thornhill Mountain.

Dermestes signatus LeC.—Scarce; a specimen compared with the type by C. A. Frost.

Dermestes talpinus Mann.

Orphilus subnitidus LeC.?

Orphilus aequalis Csy.?

BYRRHIDAE

Tylicus subcanus LeC.

Listemus acuminatus Mann. — By sifting moss.

Simpliocaria tessellata LeC.

Cytilus alternatus Say.—Along roadsides, and on flats.

Byrrhus concolor Kby.—May, 1920.

Byrrhus difficilis Csy. — From Thornhill Mountain.

Byrrhus americanus LeC. — Found among grass roots, and on flats.

Byrrhus cyclophorus Kby.—From Thornhill Mountain.

Curimopsis setulosa Mann.—Vidit H. C. Fall.

Lioligus striolatus Csy.—Vidit H. C. Fall.

OSTOMIDAE

Ostoma pippingskoeldi Mann. — Rare; on pine, and from Thornhill Mountain.

Ostoma nigrina Csy.—December 10, 1920, on rotten wood, spruce and hemlock. Rare.

Ostoma columbiana Csy.

NITIDULIDAE

Cercus pennatus Murr. — Scarce on elder bark and bush; vidit Parsons.

Colopterus truncatus Rand.—Under edges of bark on freshly cut brush, June, 1931.

Nitidula nigra Schfr. — A subspecies of *rufipes* L.; teste Parsons.

Nitidula rufipes L.—Rare under edges of bark on freshly cut brush, June, 1931.

Omosita discoidea Fab.—Frequent on bones, March 13 to June; vidit C. T. Parsons.

Omosita colon L.

Epuraea planulata Er.—(All species of this genus vidit Parsons.)

Epuraea truncatella Mann.

Epuraea spp.—Two new species, one near *avara* Rand.

Epuraea terminalis Mann.

Epuraea adumbrata Mann.

Epuraea integra Horn.

Cryptarcha ampla Er.

Glischrochilus quadrisignatus Say.—Found on stale bread.

Glischrochilus vittatus Say. — Vidit C. T. Parsons.
Glischrochilus moratus Brown.—Vidit C. T. Parsons.

RHIZOPHAGIDAE

Rhizophagus sculpturatus Mann. — Vidit H. C. Fall.
Rhizophagus remotus Lec.—Rare; in sap on cedar stump.
Rhizophagus dimidiatus Mann.—Vidit H. C. Fall.
Rhizophagus brunneus Horn.—Vidit H. C. Fall.

MONOTOMIDAE

Monotoma picipes Herbst.—Taken on a barn window in July, 1931.

CUCUJIDAE

Oryzaephilus surinamensis L. — Beneath bark.
Cathartus sp.—An unknown species.
Cathartus advena Waltl.—Det. C. A. Frost.
Pediacus fuscus Er.
Pediacus depressus Hbst.—Rare, under hemlock bark.
Cucujus clavipes var. *punicus* Mann.—Under bark.
Dendrophagus glaber LeC.—Frequent under loose bark.

EROTYLIDAE

Triplax thoracica Say.
Triplax californica LeC.—Vidit H. C. Fall.

DERODONTIDAE

Derodontus trisignatus Mann. — On white fungus in November.

CRYPTOPHAGIDAE

Salebium octodentatus Makl.
Cryptophagus lcpidus Csy.—Taken by sifting.
Cryptophagus bidentatus Makl.
Cryptophagus cellaris Scop.—March 6, 1937; vidit A. S. Nicolay.
Cryptophagus sp.—An unknown species, on cabbage in the cellar, March 18, 1937.
Cryptophagus brevipilis Csy., or near.
Cryptophagus spp.—Two unknown species; vidit H. C. Fall. One occurred on cabbage in the cellar, March 8, 1937.
Cryptophagus saginatus Sturm., or near.
Henoticus serratus Gyll.
Caenoseclis ferruginea Sahlb.—Taken when sifting fermenting grass in June and July, 1931; vidit H. C. Fall.
Atomaria fallax Csy. (In the catalogue as *Anchicera fallax* Csy.—Eds.)
Atomaria sp.—An unknown species, taken in June, 1931.
Atomaria vespertina Makl.—Vidit H. C. Fall.
Anchicera n. sp.—Taken on February 4 from a squirrel dump. Now in C. A. Frost's collection.
Anchicera ephippiata Zimm.
Anchicera ochracea var. *pennsylvanica* Csy.—Compared with type by Nicolay.

MYCETOPHAGIDAE

Mycetophagus californicus Horn.—Taken in 1933, and on Thornhill Mountain.

LATHRIDIIDAE

Lathridius costicollis LeC.
Lathridius liratus LeC.
Lathridius tardarius Deg.—In flight at windows.
Coninomus constrictus Gyll.—November 3, 1923; also by sifting, June and July, 1931.
Enicmus tenuicornis LeC.
Enicmus protensicollis Mann.—On cabbage in cellar, March 17, 1934; vidit H. C. Fall.
Enicmus fictus Fall.—On cabbage in cellar, March 6, 1937; vidit H. C. Fall.
Enicmus cordatus Bel.—June, 1931, rare.
Enicmus minutus L.—Taken while sifting, June and July, 1931; also on cabbage in cellar, March 18, 1937; vidit H. C. Fall.
Enicmus suspectus Fall.—Sifted from fermenting grass, 1937.
Corticaria ferruginea Marsh.—Sifted from moss.
Corticaria sp.—An unknown species, but "not *similata* Gyll.;" vidit H. C. Fall.
Corticaria dentigera LeC.—"columbia Fall a possibility"; vidit H. C. Fall.
Corticaria serrata Payk.
Melanophthalma sp.—An unidentified species.
Melanophthalma gibbosa Hbst.—A female.
Melanophthalma americana Mann.—Vidit H. C. Fall.
Melanophthalma cavicollis Mann.
Melanophthalma sp., near *distinguenda* Com.?

ENDOMYCHIDAE

Mycetina idahoensis Fall.—Taken from birch root, September, 1920, and June, 1931.

COCCINELLIDAE

Hyperaspis dissoluta Cr.—Rare; taken in May.
Microweisea miscella LeC.?—On poplar.
Microweisea marginata LeC.—Taken in 1933 and 1934; det. H. C. Fall.
Scymnus n. spp.—Two new species.
Scymnus marginicollis Man..
Scymnus monticola Csy.—Taken while sifting, June and July, 1931; det. Wilson.
Scymnus sp. near *nanus* Csy.—Fall says eastern and western specimens are probably different.
Psyllobora viginti-maculata Say.
Hippodamia tredecimpunctata tibialis Say.—Scarce, on alder.
Hippodamia glacialis var. *lecontei* Muls.—On everlasting flowers.
Coccinella perplexa Muls.—On wild everlasting flowers.
Coccinella transversoguttata Fald.
Coccinella transversoguttata var. *nugatoria* Muls.
Coccinella transversoguttata var. *californica* Mann.
Coccinella monticola Muls.
Coccinella perplexa var. *juliana* Muls.
Adalia bipunctata var. *quadrifasciata* Scop. According to C. Schaeffer.
Adalia frigida Schn., near var. *disjuncta* Rand.
Adalia annectans Cr., var.
Neomysia subvittata Muls.