

**A NEW HOST-PLANT IN B.C. FOR *RHOPALOSIPHUM NYMPHAEAE*
(HOMOPTERA:APHIDIDAE)**

I.V. MACRAE AND N.N. WINCHESTER
DEPT. OF BIOLOGY, UNIVERSITY OF VICTORIA
VICTORIA, B.C. V8W 2Y2

In June, 1986, while sampling insects from Eurasian watermilfoil, *Myriophyllum spicatum*, to evaluate possible biocontrol agents, adult *Rhopalosiphum nymphaeae* (L.) (Homoptera: Aphididae) were recovered from submerged plants in Ellison Lake, Kelowna, B.C. These insects were well established on the plant with probosces inserted into the stem. Individuals were observed daily for two weeks, during which reproduction occurred although the adults remained submerged, cradling bubbles of air between their legs.

R. nymphaeae were also found in samples of *M. spicatum* taken from Long Lake in Nanaimo, B.C. These insects were placed into an aquarium and fed fresh *M. spicatum* at irregular intervals. During a two month period they reproduced and alternated between feeding on the submerged plants and spending time on the surface. *R. nymphaeae* may be draining air from the plant's lacunal spaces, located throughout the stem, in much the same manner as that reported for an aquatic weevil, *Litodactylus leucogaster* (Buckingham *et al.*, 1981).

This is the first report of *R. nymphaeae* from *M. spicatum* in B.C. although it has been reported from two other submerged aquatic plants, *Callitriche stagnalis* and *Elodea canadensis* (Forbes and Chan, 1987). This species has been reported from *M. spicatum* in the southern United States (Balciunas, 1982).

R. nymphaeae has been suggested as a possible biocontrol for aquatic weeds in the past (John and Nair, 1983). This species, however, has a broad host range (Sarup *et al.*, 1973), is not an obligate aquatic, and has been reported as vectoring at least one commercially important mosaic virus from aquatic to terrestrial plants (Wyman *et al.*, 1979). For these reasons, *R. nymphaeae* may not be suitable as a biocontrol agent

ACKNOWLEDGEMENTS

We thank Dr. R. Footitt of B.R.I. for identification of specimens.

REFERENCES

- Balciunas, J.K. 1982. Insects and Other Macroinvertebrates Associated With Eurasian Watermilfoil in the United States. Tech. Rep. A-82-5. U.S. Army Engineers Waterways Expt. Stat., C.E., Vicksburg, Miss. 94 pp.
- Buckingham, G.R., Bennett, C.A., and B.M. Ross. 1981. Investigation of Two Insect Species for Control of Eurasian Watermilfoil. Tech. Rep. A-81-4. U.S. Army Engineers Waterways Expt. Stat., C.E., Vicksburg, Miss.
- Forbes, A.R., and C.K. Chan. 1987. The aphids (Homoptera: Aphididae) of British Columbia 17. A revised host plant catalogue. J. entomol Soc. Brit. Columbia 84: 72-100.
- John, K.C., and N.B. Nair. 1983. *Rhopalosiphum nymphaeae* (Homoptera: Aphididae), a control agent for *Salvinia molesta*. Entomol. Mon. 7(3): 381-384.
- Sarup, P., Singh, D.S., and R. Lal. 1973. Relative resistance of various aphids infesting terrestrial and aquatic plants to some important pesticides. Indian J. Entomol. 33: 131-135.
- Wyman, J.A., Toscano, N.C., Kido, K., Johnson, H., and K.S. Mayberry. 1979. Effects of mulching on the spread of aphid transmitted watermelon mosaic virus to summer squash. J. Econ. Entomol. 72: 139-143.